arXiv:1001.1233v2 [math.CV] 17 Mar 2010

On a Carleman formula for lunes

D. Fedchenko[1]

Abstract

In this paper we consider a simple formula for analytic continuation in a domain $D \subset \mathbb{C}$ of special form.

Introduction

Integral representations of holomorphic functions solve the classical problem on restoring a holomorphic function in a domain D by its values on ∂D. Connected with this problem, there is another one: restore holomorphic function in D by its values on a set $\Gamma \subset \partial D$.

The first result of such type was obtained by Carleman [C] for a plain domain D of special form. His idea of using *extinguishing* functions was developed in the article of Goluzin and Krylov [GK] and by Fok and Kuni [FK] for simply connected plain domains of special form. This method provides an *extinguishing* function for any subset Γ of ∂D of positive measure. Another method was proposed by Lavrent'ev in 1956 [LRS].

All these formulas and some their applications can be found in the book of Aizenberg [A].

In this notice we show a simple trick to construct Carleman formula for lunes.

1 Analytic continuation

Let $\mathbb{C}$ be the complex plane of $z = x + \imath y$, where $x, y \in \mathbb{R}$. For an open set D, denote by $\mathcal{O}(D)$ the space of functions holomorphic in D. Write $\overline{\partial}$ for the Cauchy-Riemann operator

$$\overline{\partial} = \frac{1}{2}\left(\frac{\partial}{\partial x} + \imath \frac{\partial}{\partial y}\right)$$

in $\mathbb{C}$.

Let Γ be a subset of ∂D. The problem of analytic continuation from Γ into D consists in the following. Given a function $u_0 \in C(\Gamma)$, find a function $u \in \mathcal{O}(D) \cap C(D \cup \Gamma)$ such that

---

[1] Institute of Mathematics, Siberian Federal University, Svobodny Prospect 79, Krasnoyarsk, 660041, Russia. E-mail: dfedchenk@gmail.com.

The research of the author was done in the framework of the Mikhail Lomonosov Fellowship which is supported by the Russian Ministry of Education and the Deutsche Forschungsgemeinschaft.

Key words: Analytic continuation, Carleman formulas.

2000 *Mathematics Subject Classification*: 35Cxx.

$$\begin{cases} \overline{\partial} u = 0 & in\ D, \\ u = u_0 & on\ \Gamma. \end{cases}$$

One easily specifies this problem within Cauchy problems for solutions of elliptic equations.

2 Carleman formula

Let Γ be a smooth curve in the unit disk $B(0,1)$ dividing $B(0,1)$ into two domains. Denote by D those of these domains which does not contain the origin. Such domains D are referred to as lunes. And let

$$\gamma(t) = (x(t), y(t)),\ t \geq 0,$$
$$\gamma(0) = 0$$

be a smooth curve with end point the origin, which lies in $B(0,1) \setminus D$ (see Fig. 1). Let Γ and γ intersect transversally at the origin. Fix now any curve γ with these properties.



Fig. 1. Example of D

Denote by $B(\gamma(t))$ the circle in $\mathbb{C}$ with center at the point $\gamma(t)$ and with radius $\operatorname{dist}(\gamma(t), \partial B(0,1))$.

Lemma 2.1. *For any $z \in D$, there exists point $\gamma(t_0)$ such that $z \in B(\gamma(t_0))$.*

Proof. Proof by contradiction: suppose there is a point $z \in D$ such that $z \notin B(\gamma(t))$ for all $t \geq 0$, then

$$\begin{cases} |z - \gamma(t)| \geq \operatorname{dist}(\gamma(t), \partial B(0,1)), \text{ for all } t \geq 0; \\ |z| < 1. \end{cases}$$

Let $t \to 0$. Then we have
$$\begin{cases} |z| \geq 1, \\ |z| < 1. \end{cases}$$

Obtained contradiction proofs this lemma. □

Take any numerical sequence $\gamma(t_N) = (x(t_N), y(t_N))$, $N \in \mathbb{N}$, $t_N \geq 0$ such that $\lim\limits_{N\to\infty} \gamma(t_N) = 0$.

Theorem 2.2. *If $u \in \mathcal{O}(D) \cap C(\overline{D})$, then the formula*

$$u(z) = \frac{1}{2\pi\imath} \lim_{N\to\infty} \int_\Gamma \frac{u(\zeta)}{\zeta - z} \left(\frac{z-\gamma(t_N)}{\zeta-\gamma(t_N)} \right)^{N+1} d\zeta,$$

holds for any point $z \in D$, where the convergence is uniform in z on compact subsets of D.

Proof. Fix $z \in D$. Choose N large enough, so that $z \in B(\gamma(t_N))$. Expand the Cauchy kernel as Laurent series in the variable ζ in the complement of the disk $B(\gamma(t_N))$ by

$$\frac{1}{\zeta - z} = \sum_{k=0}^{\infty} \frac{(z-\gamma(t_N))^k}{(\zeta-\gamma(t_N))^{k+1}}.$$

Consider the sequence of kernels

$$\mathfrak{C}_N(\zeta, z) = \frac{1}{2\pi\imath} \left(\frac{1}{\zeta - z} - \sum_{k=0}^{N} \frac{(z-\gamma(t_N))^k}{(\zeta-\gamma(t_N))^{k+1}} \right) \tag{2.1}$$

which we call Carleman kernel.

Using the geometric sum formula, we make the following transformation of Carleman kernels (2.1)

$$\mathfrak{C}_N(\zeta, z) = \frac{1}{2\pi\imath} \frac{1}{\zeta - z} \left(\frac{z-\gamma(t_N)}{\zeta-\gamma(t_N)} \right)^{N+1}.$$

Let $N \to \infty$. Write

$$\frac{1}{2\pi\imath} \lim_{N\to\infty} \left[\int_{\partial D} \frac{u(\zeta)}{\zeta - z} \left(\frac{z-\gamma(t_N)}{\zeta-\gamma(t_N)} \right)^{N+1} d\zeta \ - \int_{\partial D \setminus \Gamma} \frac{u(\zeta)}{\zeta - z} \left(\frac{z-\gamma(t_N)}{\zeta-\gamma(t_N)} \right)^{N+1} d\zeta \right] = \frac{1}{2\pi\imath} \lim_{N\to\infty} \int_\Gamma \frac{u(\zeta)}{\zeta - z} \left(\frac{z-\gamma(t_N)}{\zeta-\gamma(t_N)} \right)^{N+1} d\zeta. \tag{2.2}$$

Fix any compact $K \subset D$. We see that the second integral from the left-hand side in (2.2) tends to zero uniformly on K, because $q_K = \max\limits_{z \in K, \zeta \in \partial D \setminus \Gamma} \frac{|z|}{|\zeta|} < 1$. And from the holomorphy of the function $u(\zeta) \left(\frac{z-\gamma(t_N)}{\zeta-\gamma(t_N)} \right)^{N+1}$, for all $N \in \mathbb{N}$, and form the Cauchy formula we conclude that the integral over ∂D equals $u(z)$ in the domain D.

We thus arrive at the desired formula

$$u(z) = \frac{1}{2\pi\imath} \lim_{N\to\infty} \int_\Gamma \frac{u(\zeta)}{\zeta - z} \left(\frac{z-\gamma(t_N)}{\zeta-\gamma(t_N)} \right)^{N+1} d\zeta.$$

□

Example 2.3. *Let* $D = \{|z| < 1, \Re z > 0\}$ *and* $\Gamma = \{|z| < 1, \Re z = 0\}$*. As* $\gamma(t_N)$ *we take the sequence* $(-1/N, 0)$*,* $N \in \mathbb{N}$*. If* $u \in \mathcal{O}(D) \cap C(\overline{D})$*, then for any point* $z \in D \cup \Gamma$ *the formula*

$$u(z) = \frac{1}{2\pi\imath} \lim_{N\to\infty} \int_\Gamma \frac{u(\zeta)}{\zeta - z} \left(\frac{Nz+1}{N\zeta+1} \right)^{N+1} d\zeta$$

holds, where the integral converges uniformly on compact subsets of D*.*

Remark 2.4. *If* $z = 0$ *does not belong to* $\overline{D}$ *then our Carleman formula recovers one of the formulas in [A].*

References

[C] T. Carleman, *Les fonctions quasianalytiques*, Paris: Gauthier-Villars. (1926).

[GK] G. Goluzin and V. Krylov, *Generalized Carleman formula and its application to analytic continuation of functions* Mat. Sb., **40**, 144 - 149 (1933).

[FK] V. Fok and F. Kuni, *On the cutting function in dispersion relations*, Dokl. Akad. Nauk SSSR **127** (1959), 1195-1198 (Russian)

[LRS] M. M. Lavrent'ev, V. G. Romanov and S. P. Shishatskii. *Ill-Posed Problems of Mathematical Physics and Analysis* [in Russian], Nauka, Moscow (1980); English transl.: Amer. Mathem. Soc., Providence Vol. 64 (1986).

[A] L. Aizenberg, *Carleman's formulas in complex analysis*, Kluwer Acad. Publ. (1993).